

Physical Therapy Prescription for:

|              |
|--------------|
| <b>label</b> |
|--------------|

**Surgery:** L R ACL reconstruction with \_\_\_\_\_ and Medial Lateral Meniscus Repair  
**Date of Surgery:**

**Treatment:** See attached ACL Rehabilitation Protocol

**Weight bearing status:** PWB (50%) w/ crutches for \_\_\_\_\_ weeks

**Brace:**

First six weeks post-op:

- ACL hinged knee brace (TROM or equivalent) for weight bearing activities:
  - Locked when ambulating, unlocked when working on ROM (only when non-weight bearing)
  - May unlock for ambulation as soon as quad control returns
- Brace range: 0 to 90 degrees
- Brace locked at 0 degrees for sleep for first 2 weeks (or longer if hasn't achieved symmetric terminal extension)

Weeks 6 to 12:


- 1-2 week transition out of brace, provided good quad control

Weeks 12 to 14:

- No brace needed
- Focus on strengthening
- HAMSTRINGS!
- If cleared by therapist, ok to start a return to run program
- Alternate 10 min brisk walk – 10 min jog – 10 min brisk walk. Can increase jog portion each week by 2-3 minutes if tolerates well

Additional Instructions:

- SEE ATTACHED PHYSICAL THERAPY PROTOCOL
- Patella mobilizations important
- Scar massage beginning 3 weeks post-op



---

Lauren H. Redler, M.D.

Date