

Physical Therapy Prescription for:

label

Diagnosis: Elbow LUCL tear

Surgical Procedure: LUCL Reconstruction

Date of Surgery:

Precautions: No elbow ROM x 2 weeks (posterior splint at 90° in full pronation x 2 weeks)

- Slow progression w/supination, elbow extension/flexion to protect LUCL
- No weight bearing (CKC) exercises for 8-12 weeks
- No excessive flexion for 8 weeks

Immediate Post-Op (0-2 weeks) – PROTECTION PHASE:

- No elbow ROM
- Wrist ROM, gentle shoulder ROM (AROM to start week 3), and gripping exercises

Weeks 3 to 6 – RANGE OF MOTION PHASE (NO SUPINATION):

- Hinged ROM brace 30-90
- Gentle PROM elbow in neutral or pronation
- Shoulder strengthening with light dumbbells (emphasize rotator cuff & periscapular)

Weeks 7 to 8 – EARLY STRENGTHENING PHASE:

- Gradual increase ROM to 30-135, goal is full ROM by 12 weeks
- Initiate Thrower's Ten Program
- Initiate shoulder external rotation strengthening

Week 10+ – ADVANCED STRENGTHENING PHASE:

- Wean brace by week 12
- Initiate eccentric elbow flexion/extension
- Continue isotonic program: forearm/wrist
- Continue shoulder program
- Initiate 2-hand plyometric drills, advance to 1-hand plyo drills as able (~2 weeks)
- When able to do 1-hand plyo drills, ok to start advanced Thrower's Ten Program
- Return to full activities ~ 4 months
- Return to play ~ 6 months

Lauren H. Redler, M.D.

Date